

Rapid and Specific Molecular Identification of Methicillin-Resistant *Staphylococcus aureus* in Endotracheal Aspirates from Mechanically Ventilated Patients

PASCAL VANNUFFEL,¹ PIERRE-FRANÇOIS LATERRE,² MICHELE BOUYER,³ JACQUES GIGI,⁴
BERNARD VANDERCAM,⁵ MARC REYNAERT,² AND JEAN-LUC GALA^{1,3*}

Laboratory of Applied Molecular Technology,¹ and Departments of Intensive Care,² Microbiology,⁴ and
Internal Medicine,⁵ St. Luc University Hospital, and Section MSW, Operational Epidemiology and
Infectious Diseases, Queen Astrid Military Hospital,³ Brussels, Belgium

Received 24 September 1997/Returned for modification 28 January 1998/Accepted 28 April 1998

Multiplex amplification of *femA* and *mecA* genetic determinants allowed an early and rapid identification of methicillin-resistant *Staphylococcus aureus* (MRSA) in endotracheal aspirates of mechanically ventilated patients. *femA* and/or *mecA* amplification and bacteriological results were concordant in 57 of 60 samples. In all three discrepant cases, complementary bacteriological tests confirmed the presence of MRSA first identified by molecular analysis. These results underline the value and rapidity of this molecular diagnosis for MRSA infection and control surveillance in intensive care units. Rapid MRSA detection is expected to have a significant clinical impact not only on patient outcome but also on the costs for isolation and treatment.

The frequencies of methicillin-resistant *Staphylococcus aureus* (MRSA) and methicillin-resistant coagulase-negative staphylococcus (MR-CNS) infections are a worldwide concern, especially in intensive care units (ICU) (4, 12, 16). Risk factors include prolonged tracheal intubation and mechanical ventilation in critically ill patients (28, 30). In these patients, MRSA-mediated nosocomial pulmonary infections are associated with a high mortality and morbidity (1, 11, 23). Mechanisms of bacterial colonization and airway inoculation include aspiration of secretions contaminated with the pathogenic organism (10). MRSA can be acquired endogenously or during hospitalization by cross-contamination from colonized health care workers or other chronically infected patients (21, 22). Therefore, rapid and specific detection of MRSA colonization in upper and lower respiratory tracts is of paramount importance for appropriate therapeutic management and patient isolation. Unfortunately, conventional MRSA diagnosis in tracheal aspirates is adversely affected by several factors: the time required for proper bacterial identification and an accurate susceptibility test, frequent colonization of the upper airways by gram-negative bacilli (6), potential growth inhibition of staphylococcal species by gram-negative bacteria (18), great variability of growth conditions for MRSA (27), and prior antibiotic treatment, which may reduce the sensitivity of microorganism identification (7).

In staphylococcal species, resistance to methicillin and other β -lactam antibiotics is primarily mediated by the overproduction of an additional altered penicillin-binding protein (PBP2a) (13). The *mecA* gene, the structural determinant encoding PBP2a, is highly conserved among the methicillin-resistant species but is absent from susceptible strains, making it a useful molecular marker of β -lactam resistance in all staphylococci (24, 25). Another chromosomal element, *femA*, which cooperates with *mecA* for the expression of β -lactam resistance, appears to be a unique feature of *S. aureus* (3, 26, 29).

We recently validated in vitro a multiplex PCR where coamplification of both determinants clearly distinguished susceptible (lacking *mecA*) from resistant (*mecA*⁺) staphylococci, as well as distinguishing *S. aureus* (*femA*⁺) from coagulase-negative staphylococci (lacking *femA*) (29) (Fig. 1). Although very attractive, such a molecular identification still awaited clinical evaluation. In this study, we prospectively assessed the value of the multiplex assay for endotracheal aspirates (ETA). After family consent, samples were collected from mechanically ventilated patients during and a few months after an outbreak of MRSA infections.

(This study was presented at the 37th Interscience Conference on Antimicrobial Agents and Chemotherapy, Toronto, Canada, October 1997 [28a].)

A total of 48 patients, treated in four contiguous ICU, were included in the study, 33 of them being assessed over a period of 3 weeks during the outbreak and 15 being assessed a few months afterward. ETA were collected as follows. Five milliliters of sterile physiological saline was first instilled via the tracheal tube. After two or three artificial respiratory cycles, up to 3 ml of fluid was gently aspirated into a sterile tube via a sterile flexible cannula connected to a vacuum pump. Sixty duplicate ETA were collected for both microbiological examination and multiplex PCR analysis. Specimens for molecular analysis were either prepared immediately or maintained for a maximum of 24 h at 4°C until processed. Conventional identification of species and susceptibility tests were performed by disk diffusion testing with oxacillin in accordance with the National Committee for Clinical Laboratory Standards criteria described previously (15, 20). For the multiplex PCR assay, clinical specimens were homogenized in 5 ml of TE buffer (20 mM Tris HCl [pH 8.0], 10 mM EDTA) containing 2% (wt/vol) sodium dodecyl sulfate. The homogenate (1.5 ml) was then centrifuged for 5 min at 7,500 $\times g$. The cellular pellet was washed once with TE buffer, lysed in the presence of 1% (vol/vol) Triton X-100 and 50 μ g of lysostaphin (Sigma Chemical Co., St. Louis, Mo.), and incubated for 15 min at 37°C. Lysis was completed by adding 100 μ g of proteinase K (Boehringer, Mannheim, Germany). The lysate was incubated for another 15 min at 55°C and 5 min at 95°C. It was centrifuged

* Corresponding author. Mailing address: Laboratory of Applied Molecular Technology, Clos Chapelle aux Champs, 30-UCL/30.46, B-1200, Brussels, Belgium. Phone: 32-2-764 3165. Fax: 32-2-764 3959. E-mail: gala@sang.ucl.ac.be.




FIG. 1. Differential identification of staphylococcal species and methicillin-resistant strains. DNA fragments resulting from the amplification of each gene were separated by electrophoresis on agarose gel. Lanes 1 and 2, amplification of *femA* and *mecA* determinants with primers F1/F2 and M1/M2, respectively; lane 3, molecular weight markers (values [in base pairs] in order from top to bottom are 1,114, 900, 692, 501/486, 404, 320, 242, and 190); lanes 4 to 7, simultaneous amplification of both markers from MRSA (lane 4; amplification of both determinants) and MSSA (lane 5; amplification of *femA* alone) DNA and from MR-CNS (lane 6; amplification of *mecA* alone) and MS-CNS (lane 7; no amplification) DNA.

at $4,000 \times g$ for 5 min. In order to purify bacterial DNA, 200 μ l of the supernatant was then filtered on a NucleoSpin C+T column (Macherey-Nagel, Düren, Germany) and eluted with 200 μ l of sterile H₂O, according to the manufacturer's protocol. Two different amounts of the DNA suspension (2 and 20 μ l) were subjected to multiplex PCR amplification as previously described (29). Primers used were 5'-TGGCTATCGTTCACAATCG-3' and 5'-CTGGAACCTTGTTGAGCAGAG-3' for *mecA* and 5'-CTTACTTACTGGCTGTACCTG-3' and 5'-ATGTCGCTTGTTATGTGC-3' for *femA*, yielding 310- and 686-bp fragments, respectively (Fig. 1). The 40 PCR cycles were carried out in a model 2400 thermocycler (Perkin-Elmer, Foster City, Calif.) as follows: denaturation at 92°C for 20 s, annealing at 58°C for 20 s, and DNA extension at 72°C for 20 s, with increments of 2 s per cycle for the denaturation and extension segments. Amplified, ethidium bromide-stained DNA fragments were visualized after electrophoresis on agarose gel.

Molecular and conventional tests were performed in different laboratories, and results were compared (Table 1). In the multiplex amplification, identical results were obtained with either 2 or 20 μ l of the DNA suspension. A perfect correlation

TABLE 1. Bacteriological and molecular results for ETA^a

No. of ETA	Identification	
	Phenotypic (bacteriological tests)	Genotypic (multiplex PCR strategy)
Correlations (<i>n</i> = 57)		
25	MRSA	<i>femA</i> ⁺ / <i>mecA</i> ⁺
10	MSSA	<i>femA</i> ⁺ /no <i>mecA</i>
6	MR-CNS	no <i>femA</i> / <i>mecA</i> ⁺
6	MS-CNS	no <i>femA</i> /no <i>mecA</i>
5	Gram-negative bacteria	no <i>femA</i> /no <i>mecA</i>
5	Normal flora	no <i>femA</i> /no <i>mecA</i>
Discrepancies (<i>n</i> = 3)		
1	Gram-negative bacteria ^b	<i>femA</i> ⁺ / <i>mecA</i> ⁺
2	No MRSA ^c	<i>femA</i> ⁺ / <i>mecA</i> ⁺

^a The 25 MRSA-positive ETA were found in 15 of 33 patients during the outbreak; the 5 ETA containing normal flora were collected from three patients. All other results correspond to one ETA per patient.

^b *P. aeruginosa* coinfection.

^c MRSA nasal carriage.

between genotypic and phenotypic analyses was found for 57 ETA. The *mecA* and *femA* determinants were amplified from every ETA containing MRSA (*n* = 25), as determined by standard bacteriological methods. Single *femA* or *mecA* signals were found in specimens containing either methicillin-susceptible *S. aureus* (MSSA) (*n* = 10) or MR-CNS (*n* = 6), respectively. On the other hand, no signal was obtained from ETA colonized with gram-negative bacteria (*n* = 5) or methicillin-susceptible CNS (MS-CNS) (*n* = 6) and from five ETA containing normal pharyngeal flora.

Of note, discrepancies were found in three cases where *femA* and *mecA* markers were both amplified in ETA but no bacteriological evidence of staphylococci was found in the corresponding culture. One of the three cultures was massively positive for *Pseudomonas aeruginosa*. However, further identification on hyperselective medium (mannitol-salt-agar), prompted by the discrepant molecular result, identified MRSA in this ETA. For the other two cases, bacteriological controls revealed MRSA nasal carriage at the time of ETA specimen collection. Successive cultures repeated over the next few days also identified MRSA in ETA.

These results extend our previous in vitro data (29). They emphasize the sensitivity and specificity of the multiplex PCR strategy for detecting MRSA in ETA. The entire procedure can be completed in less than 6 h, either on the day of sample collection or the next day. This is quicker than conventional identification and susceptibility tests (48 to 72 h). Interestingly, the current data suggest that MRSA molecular detection is valuable for samples coinfecting by fast-growing gram-negative bacteria such as *P. aeruginosa*, a potential cause of false-negative results by standard methods. Such a rapid and accurate MRSA identification may be applied to other sites such as nares, sputum, and wounds.

Accordingly, we can expect this method to have a positive economic impact. Effective but expensive barrier isolations are indeed recommended to control MRSA, especially for patients at high risk for MRSA carriage (transfer from a nursing home or other chronic care facilities and hospitals) (2, 5, 14). On the other hand, empiric glycopeptide treatment has been officially recommended for nosocomial pneumonia with risk factors (2), because delayed administration of adequate antibiotic therapy is associated with a greater risk of hospital mortality (reviewed in reference 17). Whereas the cost per sample of a conventional culture appears to be in the same range as that for our molecular assay (14) (data not shown), rapid same-day bacterial identification and susceptibility testing by a molecular method would prevent unnecessary isolation procedures and inappropriate empiric glycopeptide treatment, both of which are responsible for high additional costs (8, 14, 19). Moreover, rapid testing has been shown to have a major impact on care and outcome (i.e., resulting in a lower mortality rate, fewer laboratory studies, and shorter stays in the ICU, etc.) (9).

The project was funded by JSM-R&D-T2, the Joint Staff section of the Belgian Army supporting research and development (grant G95/01). P.V. was supported in part by a grant from Eli Lilly Benelux.

We thank J. Lünzer (Macherey-Nagel) for the generous gift of NucleoSpin C+T kits and Anne Vandenberghe for so enthusiastically and carefully collecting the endotracheal samples. We also thank M. Philippe, Département of Clinical Chemistry, for stimulating discussions and providing laboratory facilities.

REFERENCES

- Al Ujayli, B., D. A. Nafziger, and L. Saravolatz. 1995. Pneumonia due to *Staphylococcus aureus* infection. Clin. Chest Med. 16:111-120.
- American Thoracic Society. 1996. Hospital-acquired pneumonia in adults: diagnosis, assessment of severity, initial antimicrobial therapy, and preven-

- tive strategies. A consensus statement. *Am. J. Respir. Crit. Care Med.* **153**: 1711–1725.
3. **Berger-Bachi, B., L. Barberis-Maino, A. Strassle, and F. H. Kayser.** 1989. *FemA*, a host-mediated factor essential for methicillin resistance in *Staphylococcus aureus*: molecular cloning and characterization. *Mol. Gen. Genet.* **219**:263–269.
 4. **Blumberg, L. H., and K. P. Klugman.** 1994. Control of methicillin-resistant *Staphylococcus aureus* bacteraemia in high-risk areas. *Eur. J. Clin. Microbiol. Infect. Dis.* **13**:82–85.
 5. **Casewell, M. W.** 1995. New threats to the control of methicillin-resistant *Staphylococcus aureus*. *J. Hosp. Infect.* **30**:465–471.
 6. **Cazzadori, A., G. Di-Perri, S. Vento, S. Bonora, D. Fendt, M. Rossi, M. Lanzafame, F. Mirandola, and E. Concia.** 1997. Etiology of pneumonia following isolated closed head injury. *Respir. Med.* **91**:193–199.
 7. **Chastre, J., J. Y. Fagon, and C. Laner.** 1992. Procedures for the diagnosis of pneumonia in ICU patients. *Intensive Care Med.* **18**:S10–S17.
 8. **Doebelling, B. N., and R. P. Wenzel.** 1990. The direct costs of universal precautions in a teaching hospital. *JAMA* **264**:2083–2087.
 9. **Doern, G. V., R. Vautour, M. Gaudet, and B. Levy.** 1994. Clinical impact of rapid in vitro susceptibility testing and bacterial identification. *J. Clin. Microbiol.* **32**:1757–1762.
 10. **Estes, R. J., and G. U. Meduri.** 1995. The pathogenesis of ventilator-associated pneumonia. I. Mechanisms of bacterial transcolonization and airway inoculation. *Intensive Care Med.* **21**:365–383.
 11. **Fagon, J. Y., J. Chastre, A. Vuagnat, J. L. Trouillet, A. Novara, and C. Gibert.** 1996. Nosocomial pneumonia and mortality among patients in intensive care units. *JAMA* **275**:866–869.
 12. **Fridkin, S. K., S. F. Welbel, and R. A. Weinstein.** 1997. Magnitude and prevention of nosocomial infections in the intensive care unit. *Infect. Dis. Clin. N. Am.* **11**:479.
 13. **Hackbart, C., and H. Chambers.** 1989. Methicillin-resistant staphylococci: genetics and mechanisms of resistance. *Antimicrob. Agents Chemother.* **33**:991–999.
 14. **Jernigan, J. A., M. A. Clemence, G. A. Stott, M. G. Titus, C. H. Alexander, C. M. Palumbo, and B. M. Farr.** 1995. Control of methicillin-resistant *Staphylococcus aureus* at a university hospital: one decade later. *Infect. Control Hosp. Epidemiol.* **146**:686–696.
 15. **Kloos, W. E., and D. W. Lambe, Jr.** 1991. *Staphylococcus*, p. 222–237. In A. Balows, W. J. Hausler, Jr., K. L. Herrmann, H. D. Isenberg, and H. J. Shadomy (ed.), *Manual of clinical microbiology*, 5th ed. American Society for Microbiology, Washington, D.C.
 16. **Kloos, W. E., and T. L. Bannerman.** 1994. Update of clinical significance of coagulase-negative staphylococci. *Clin. Microbiol. Rev.* **7**:117–140.
 17. **Kollef, M. H.** 1998. The importance of initial antibiotic selection in ventilator-associated pneumonia, p. 300–309. In J. L. Vincent (ed.), *Yearbook of intensive care and emergency medicine 1998*. Springer-Verlag, Berlin, Germany.
 18. **Machan, Z. A., T. L. Pitt, W. White, D. Watson, G. W. Taylor, P. J. Cole, and R. Wilson.** 1991. Interaction between *Pseudomonas aeruginosa* and *Staphylococcus aureus*: description of an anti-staphylococcal substance. *J. Med. Microbiol.* **34**:213–217.
 19. **Marshall, J. C., and D. C. Evans.** 1998. Antimicrobial therapy for ICU-acquired infection: time for reappraisal, p. 283–291. In J. L. Vincent (ed.), *Yearbook of intensive care and emergency medicine 1998*. Springer-Verlag, Berlin, Germany.
 20. **National Committee for Clinical Laboratory Standards.** 1994. Performance standards for antimicrobial disk susceptibility tests, 5th ed. M2-A5. National Committee for Clinical Laboratory Standards, Villanova, Pa.
 21. **Phillips, L. G., J. P. Heggers, and M. C. Robson.** 1992. Burn and trauma units as source of methicillin-resistant *Staphylococcus aureus*. *Burn Care Rehabil.* **13**:293–297.
 22. **Pujol, M., C. Pena, R. Pallares, J. Ariza, J. Ayats, M. A. Dominguez, and F. Gudiol.** 1996. Nosocomial *Staphylococcus aureus* bacteremia among nasal carriers of methicillin-resistant and methicillin-susceptible strains. *Am. J. Med.* **100**:509–516.
 23. **Rello, J., A. Torres, M. Riccart, J. Valles, J. Gonzalez, A. Artigas, and R. Rodriguez-Roisin.** 1994. Ventilator-associated pneumonia by *Staphylococcus aureus*. Comparison of methicillin-resistant and methicillin-sensitive episodes. *Am. J. Respir. Crit. Care Med.* **150**:1545–1549.
 24. **Ryffel, C., W. Tesch, I. Birch-Machin, P. E. Reynolds, L. Barberis-Maino, F. H. Kayser, and B. Berger-Bachi.** 1990. Sequence comparison of *mecA* genes isolated from methicillin-resistant *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Gene* **94**:137–138.
 25. **Suzuki, E., K. Hiramatsu, and T. Yokota.** 1992. Survey of methicillin-resistant clinical strains of coagulase-negative staphylococci for *mecA* gene distribution. *Antimicrob. Agents Chemother.* **36**:429–434.
 26. **Unal, S., J. Hoskins, J. E. Flokowitsch, C. Y. E. Wu, D. A. Preston, and P. L. Skatrud.** 1992. Detection of methicillin-resistant staphylococci by using polymerase chain reaction. *J. Clin. Microbiol.* **30**:1685–1691.
 27. **Unal, S., K. Werner, P. DeGirolami, F. Barsanti, and G. Eliopoulos.** 1994. Comparison of tests for detection of methicillin-resistant *Staphylococcus aureus* in a clinical microbiology laboratory. *Antimicrob. Agents Chemother.* **38**:345–347.
 28. **Valles, L., C. Leon, and F. Alvarez-Lerma.** 1997. Nosocomial bacteremia in critically ill patients: a multicenter study evaluating epidemiology and prognosis. *Clin. Infect. Dis.* **24**:387–395.
 - 28a. **Vannuffel, P., M. Bouyer, P.-F. Laterre, J. Gigi, A. Vandenberghe, B. Vandercam, M. Reynaert, and J.-L. Gala.** 1997. Rapid and specific identification of MRSA in endotracheal aspirates from mechanically ventilated ICU patients, abstr. D-25, p. 87. In *Abstracts of the 37th Interscience Conference on Antimicrobial Agents and Chemotherapy*. American Society for Microbiology, Washington, D.C.
 29. **Vannuffel, P., J. Gigi, H. Ezzedine, B. Vandercam, M. Delmee, G. Wauters, and J.-L. Gala.** 1995. Specific detection of methicillin-resistant *Staphylococcus* species by multiplex PCR. *J. Clin. Microbiol.* **33**:2864–2867.
 30. **Vincent, J. L., D. J. Bihari, P. M. Suter, H. A. Bruining, J. White, M. H. Nicolas-Chanoine, M. Wolff, R. C. Spencer, and M. Hemmer.** 1995. The prevalence of nosocomial infection in intensive care units in Europe. Results of the European Prevalence of Infection in Intensive Care (EPIC) Study. EPIC International Advisory Committee. *JAMA* **274**:639–644.